

Notes and Comments

***Hexacladia smithii* Ashmead, 1891 (Hymenoptera: Encyrtidae) in Panama: first record and new association with *Berecynthus hastator* (Fabricius, 1798) (Hemiptera: Pentatomidae)**

B. A. Zachrisson^a , J. Grazia^b , V. A. Costa^c , G. Gutierrez^a 

^a Instituto de Innovación Agropecuaria de Panamá – IDIAP, Centro de Innovación Agropecuaria Oriental – CIAOr, Laboratorio de Entomología Agrícola – Control Biológico de Plagas, Chepo, Panamá

^b Universidade Federal do Rio Grande do Sul – UFRGS, Departamento de Zoologia, Programa de Pós-Graduação em Biologia Animal, Porto Alegre, RS, Brasil

^c Instituto Biológico, Centro Avançado de Pesquisa e Desenvolvimento em Sanidade Agropecuária, Campinas, SP, Brasil

Berecynthus hastator (Fabricius, 1798) is synonym of *Berecynthus delirator* (Fabricius, 1787) (Grazia and Hildebrandt, 1982; Rider and Rolston, 1995), whose geographical distribution is limited to the neotropics. Currently, there are records of this species in Bolivia, Brazil, Costa Rica, Colombia, French Guyana, Mexico, Nicaragua, Peru, Puerto Rico, Suriname, Venezuela and Panama (Grazia et al., 2015; Segarra-Carmona et al., 2015; Cazorla and Moreno, 2020; Rider, 2021; Dellapé, 2021; GBIF, 2022). In Panama *B. hastator* was reported in Coco Solo, Province of Colon (09° 22' 21.43" N, 79° 52' 52.32" W), Puerto Armuelles (08° 39' 09" N, 82° 07' 24" W), Bayano (09° 06' 58" N, 78° 46' 24" W), Kobe (08° 55' N, 79° 34' W), Chorrera (08° 52' 49" N, 79° 47' W), Santiago (08° 07' N, 80° 59' W), Las Cumbres (09° 06' N, 79° 32' W) (Grazia and Hildebrandt, 1982; GBIF, 2022). Adults of *B. hastator* were collected in Panama with light traps and its association with agricultural crops and uncultivated plants was not confirmed (GBIF, 2022); the voucher specimens were deposited in the collection of Texas A&M University, College Station, Texas, USA; Illinois Natural History Survey Insect Collection, Illinois, USA, and Smithsonian Tropical Research Institute, Panama (GBIF, 2022). The adults of *B. hastator* collected in Panama with of light traps and its association with agricultural crops and uncultivated plants were not confirmed (GBIF, 2022). The impact of *B. hastator* on crops of economic importance is limited; this insect is mentioned as a possible vector of *Phytomonas staheli* McGhee and McGhee, 1979 (Trypanosomatidae), a possible etiological agent of the “Marchitez Sorpresiva” in species of Arecaceae, among these *Cocos nucifera* L., *Elaeis guineensis* Jack (Segeren, 1982). In addition, the associations between *B. hastator* and *Gossypium hirsutum* L. and *Phaseolus* sp. (Tellez and Maes, 1991), *Oryza sativa* L. (Krinski, 2014; Rider, 2021), *Rottboellia cochinchinensis* (Lour.) Clayton (Haywood et al., 1994) and *Sabal mexicana* Mart. (Brailovsky et al., 1992) were confirmed.

Hexacladia smithii Ashmead, 1891 is a gregarious endoparasitoid present in the neotropics which develops in nymphs and adults of some species of Coreidae (Baldin et al., 2010), Pentatomidae (Santis, 1979; Corrêa-Ferreira et al., 1998) and Scutelleridae (Santis, 1980). Records of *H. smithii* were confirmed in Brazil (Ashmead, 1891), Costa Rica (Noyes, 2010), Nicaragua (Noyes, 2010), Puerto Rico (Noyes, 2010), and Trinidad and Tobago (Burks, 1972; Noyes, 2010). Some species of economic importance such as *Arvellius albopunctatus* (DeGeer, 1773) (Panizzi and Silva, 2010), *Edessa meditabunda* (Fabricius, 1794) (Cuezzo and Fidalgo, 1997), *Euschistus heros* (Fabricius, 1798) (Corrêa-Ferreira et al., 1998) and *Diceraeus furcatus* (Fabricius, 1775) (cited as *Dichelops furcatus*, Panizzi and Silva, 2010), were parasitized by *H. smithii*, in the soybean crops (*Glycine max* L.) and tomato (*Solanum lycopersicum* L.) (Panizzi and Silva, 2010), among others. In Panama, there is limited information on new records of parasitoids and insects-pest of economic importance and the trophic associations they may present. Therefore, the confirmation of the biotic relationship between *H. smithii* and *B. hastator* is considered a relevant contribution to strengthening biological control and integrated pest management programs in emerging insects-pest.

Adults of *B. hastator* were randomly collected in 25 tomato plants (*S. lycopersicum* - Cultivar “IDIAP T-7”) by means of insect aspirators adapted for the capture of parasitoids, during the periods of February 2017, March 2018 and February 2019. The experimental tomato plots of 50 m² were cultivated in a family farming system located in Cabra, Pacora, Panama (09° 07' N, 79° 19' W). Dr. Jocelia Grazia (Universidade Federal do Rio Grande do Sul) made the identification and confirmation of *B. hastator*. The diagnosis of this species indicates that the genus *Berecynthus* is close to *Dichelops* Spinola, *Padaeus* Stål, *Euschistus* Dallas (Grazia, 1978) and more recently to *Diceraeus* Dallas (Barão et al., 2020). *Berecynthus hastator* (Fabricius, 1798) is the only valid species of the genus, characterized by the clypeus

^a e-mail: bazsalam@gmail.com

Received: April 18, 2022 – Accepted: May 30, 2022



This is an Open Access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

longer than mandibular plates, differing from the other closer genera, except *Euschistus*, that includes some species with clypeus longer. The main differences of *Berecyntus* are found in the morphology of genitalia for both sexes (Figure 1). Voucher specimens collected in 2017 (3♀, 2♂) and 2018 (2♀, 1♂), were deposited in the collections of the Museum of Invertebrates G.B. Fairchild of University of Panama, Panama and the Laboratory of Entomological Systematics (LES) of the Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil (1♀, 2♂).

Adults of *B. hastator* collected in the experimental plots were separated and transferred to transparent plastic containers (7 cm x 5 cm x 5 cm) with a moistened paper towel at the base of the container and covered with a




Figure 1. *Berecyntus hastator* (Fabricius, 1798) (Hemiptera: Pentatomidae). (a) body, dorsal view; (b) body, ventral view; (c) female genital plate. Photo: Dr. Lurdiana D. Barros.



Figure 2. *Hexacladia smithii* (Ashmead, 1891) (Hymenoptera: Encyrtidae). (a) body, lateral view; (b) male fore wing; (c) female fore wing. Photo: Dr. Valmir A. Costa.

tulle fabric mesh. Individuals were separated into groups of approximately 10 adults of *B. hastator* per container, where they were fed peanuts, bean pods and a 90% honey solution, offered in small swabs of cotton. In order to maintain temperature, relative humidity, and constant photoperiod, the insects were transferred to regulated chambers at 28+2 °C, 80% of relative humidity and 12 hours of photophase, located in the laboratory of Biological Pest Control of the Instituto de Innovación Agropecuaria de Panamá (IDIAP), in Chepo, Panama. Adults of *H. smithii* emerged from *B. hastator* were identified by Dr. Valmir Costa (Instituto Biológico, Campinas, SP, Brazil) following the key of Noyes (2010). This endoparasitoid presents reddish-brown color (Figure 2a) and the basitarsus of mid legs, in both sexes, a line of about 10 stout pegs ventrally. The male's fore wing, post marginal vein at least 1.5 x as long as stigmal vein (Figure 2b), and the female's fore wing is characterized by the subapical infuscate area broadly connected to median infuscate band, costal cell with less than 10 setae dorsally and at least 2 or 3 setae ventrally, and postmarginal vein clearly longer than stigmal vein (Figure 2 c). Voucher specimens (5♀, 3♂) were deposited in the collections of the Museum of Invertebrates G.B. Fairchild of the University of Panama, Panama, and the Oscar Monte Collection of Entomophagous Insects (3♀, 4♂) (IBCBE-007229-007235) of the Instituto Biológico, Campinas, São Paulo, Brazil.

The circular holes caused by the emergence of adults of *H. smithii* in the abdomen of adults of *B. hastator* and the quantification of emerged adults of *H. smithii* allowed to calculate the rate of (%) natural parasitism.

$$\% \text{ Parasitism} = [(\text{Total Parasitized Adults} / \text{Total Collected Adults}) 100]$$

The natural parasitism of *H. smithii* in adults of *B. hastator*, over the years 2017, 2018 and 2019, was 39.13% (9/23), 56.25% (9/16), 38.09% (8/21), respectively. Therefore, the first report of *H. smithii* is considered a new trophic association between this endoparasitoid and *B. hastator*, for Panama. No damage caused by adults of *B. hastator* were observed on tomato plants. The data submitted by GBIF (2022) indicated that in Panama the collection of *B. hastator* were made by means of light traps, thus the trophic association "S. lycopersicum - *B. hastator*" was not confirmed. Noyes (2019) confirmed that there were no previous records of *H. smithii* in Panama, information that was validated by GBIF (2022). The relevance of the information presented strengthens the Pentatomidae species management programs, based on the adaptation and colonization of *H. smithii*, in the agroecological zones of Panama.

Acknowledgments

B. Zachrisson (corresponding author) thanks the Sistema Nacional de Investigación (SNI) of the Secretaría Nacional de Ciencia, Tecnología e Innovación (SENACYT) of Panama for the grant (SNI-19-2020) and the financial support, allowing the conduction of this research. V.A. Costa thanks the INCT-Hympar (Instituto Nacional de Ciência e Tecnologia dos Hymenoptera Parasitoides, CNPq Process

465562/2014-0) for financial support. J. Grazia thanks to CNPq for the fellowship grant (PQ #310747/2020-1) and to L.D. Barros for the photos of *B. hastator*.

References

- ASHMEAD, W.H., 1891. An encyrtid with six-branched antennae. *Insect Life*, vol. 3, pp. 455-457.
- BALDIN, E.L.L., FUJIHARA, R.T., BOIÇA JUNIOR, A.L. and ALMEIDA, M.C., 2010. Parasitismo de percevejos-praga do maracujazeiro no Brasil por *Hexacladia smithii* Ashmead (Hymenoptera: encyrtidae). *Neotropical Entomology*, vol. 39, no. 2, pp. 306-307. <http://dx.doi.org/10.1590/S1519-566X2010000200025>. PMID:20498972.
- BARÃO, K.R., FERRARI, A. and GRAZIA, J., 2020. Phylogenetic analysis of the *Euschistus* group (Hemiptera: Pentatomidae) suggests polyphyly of *Dichelops Spinola, 1837* with the erection of *Diceraeus* Dallas, 1851, stat. rev. *Austral Entomology*, vol. 59, no. 4, pp. 770-783. <http://dx.doi.org/10.1111/aen.12489>.
- BRAILOVSKY, H., CERVANTES, L. and MAYORGA, C., 1992. *Hemiptera: Heteroptera de México XLIV. Biología, Estadios Ninfales y Fenología de la tribu Pentatomini (Pentatomidae) en la Estación de Biología Tropical "Los Tuxtlas"*. Veracruz: Universidad Nacional Autónoma de México, 204 p.
- BURKS, B.D., 1972. The genus *Hexacladia* Ashmead (Hymenoptera: encyrtidae). *Proceedings of the Entomological Society of Washington*, vol. 74, no. 4, pp. 363-371.
- CAZORLA, D. and MORENO, O.M., 2020. Nuevos registros de Pentatomidae (Hemiptera: Heteroptera) en el Estado de Falcón, Venezuela. *Revista Nicaraguense Entomológica*, vol. 197, pp. 3-54.
- CORRÊA-FERREIRA, B.S., NUNES, M.C. and UGUCCIONI, L.D., 1998. Occurrence of *Hexacladia smithii* Ashmead parasitoid of *Euschistus heros* (F.) adults in Brazil. *Anais da Sociedade Entomológica do Brasil*, vol. 27, no. 3, pp. 495-498. <http://dx.doi.org/10.1590/S0301-80591998000300022>.
- CUEZZO, F. and FIDALGO, P., 1997. *Hexacladia smithii* Ashmead (Hymenoptera: Encyrtidae): a new record for Argentina and two new pentatomid hosts recorded, *Antiteuchus variolosus* Westwood and *Edessa meditabunda* F. (Hemiptera: Pentatomidae). Entomologist, vol. 116, no. 1, pp. 11-14.
- DELLAPÉ, G., 2021. New records of neotropical Pentatominae (Heteroptera: Pentatomidae). *Entomological Communications*, vol. 3, ec03004.
- SANTIS, L., 1979. *Catálogo de los himénopteros calcidoideos de América al sur de los Estados Unidos*. La Plata: Comisión de Investigaciones Científicas, 488 p.
- SANTIS, L., 1980. *Catálogo de los himenópteros brasileiros de la serie parasitica incluyendo Bethylloidea*. Curitiba: Editora da Universidade Federal do Paraná, 187 p.
- GLOBAL BIODIVERSITY INFORMATION FACILITY – GBIF, 2022 [viewed 30 January 2022]. *Spermacoce verticillata* L.. Available from: <https://www.gbif.org/species/2918574>.
- GRAZIA, J., 1978. Revisão do gênero *Dichelops* Spinola, 1837 (Heteroptera, Pentatomidae, Pentatomini). *Iheringia. Série Zoologia*, vol. 53, pp. 3-119.
- GRAZIA, J. and HILDEBRANT, R., 1982. Revisão do gênero *Berecythus* Stål, 1862 (Heteroptera, Pentatomidae, Pentatomini). *Revista Brasileira de Entomologia*, vol. 26, no. 2, pp. 173-182.
- GRAZIA, J., PANIZZI, A.R., GREVE, C., SCHWERTNER, C.F., CAMPOS, L.A., GABERLOTTO, T.A. and FERNANDES, J.A.M., 2015. Stink bugs (Pentatomidae). In: A.R. PANIZZI and J. GRAZIA, eds. *True bugs (Heteroptera) of the Neotropics*. Dordrecht: Springer, pp. 681-756. http://dx.doi.org/10.1007/978-94-017-9861-7_22.
- HAYWOOD, P.J., LANCASTERZS, B. and EASTWOOD, D., 1994. Insects associated with, and some diseases of *Rottboellia cochinchinensis* (Lour.) W.D. Clayton in Guyana. In: *Proceedings of 25th Conference, West Indies Sugar Technologists*, April 18-21, Belize. Belize: West Indies Sugar Technologists. pp. 101-111.
- KRINSKI, D., 2014. Artrópodes associados ao arroz de terras altas, *Oryza sativa*, em Novo Progresso, Estado do Pará: níveis de danos e estratégias para manejo. Paraná: Universidade Federal do Paraná, 326 p. Tese de Doutorado em Zoologia.
- NOYES, J.S., 2010. Encyrtidae of Costa Rica (Hymenoptera: Chalcidoidea), 3. Subfamily Encyrtinae: Encyrtini, Echthrophlexiellini, Discodini, Oobiini and Ixodiphagini, parasitoids associated with bugs (Hemiptera), insect eggs (Hemiptera, Lepidoptera, Coleoptera, Neuroptera) and ticks (Acari). *Memoirs of the American Entomological Institute*, vol. 84, pp. 1-848.
- NOYES, J.S., 2019 [viewed 15 April 2022]. *Universal Chalcidoidea Database*, 2019 [online]. Available from: <https://www.nhm.ac.uk/our-science/data/chalcidooids/database/>.
- PANIZZI, A.R. and SILVA, J.J., 2010. New records of pentatomids as hosts of *Hexacladia smithii* Ashmead (Hymenoptera: Encyrtidae) in southern Brazil. *Neotropical Entomology*, vol. 39, no. 4, pp. 678-679. <http://dx.doi.org/10.1590/S1519-566X2010000400034>. PMID:20878011.
- RIDER, D.A. and ROLSTON, L.H., 1995. Nomenclatural changes in the Pentatomidae (Hemiptera: pentatomidae). *Proceedings of the Entomological Society of Washington*, vol. 97, pp. 845-855.
- RIDER, D.A., 2021 [viewed 20 December 2021]. *Pentatomidea Home Page* [online]. Available from: www.ndsu.edu/pubweb/~rider/Pentatomidea/index.htm.
- SEGEREN, P., 1982 [viewed 20 December 2021]. Preliminary study on the vectors of "Harrtrot" disease of coconut in Suriname. *Surinaamse Landbouw* [online], vol. 30, no. 1, pp. 17-23. Available from: <https://www.cabdirect.org/cabdirect/abstract/19826741362>
- SEGARRA-CARMONA, A.E., FRANQUI, R.A., PEREZ-MARTÍNEZ, H. and MORALES-GRAJALES, A., 2015. New records for stink bugs (Heteroptera: Pentatomidae) in Puerto Rico. *The Journal of Agriculture of the University of Puerto Rico*, vol. 99, no. 2, pp. 167-178. <http://dx.doi.org/10.46429/aupr.v99i2>.
- TELLEZ, J. and MAES, J.M., 1991. Policultivo maíz- frijol como elemento de lucha biológica contra *Dalbulus maidis* (Homoptera: Cicadellidae). II. Influencia sobre plagas de maíz y frijol. *Revista Nicaraguense Entomológica*, vol. 16, pp. 9-20.